

# Differential Diagnosis of Metatarsus Adductus

The author clarifies how to make the correct diagnosis of this condition.

BY MARC A. BENARD, DPM



## Goals and Objectives

- 1) To improve the practitioner's ability to assess metatarsus adductus
- 2) To improve the practitioner's ability to differentiate metatarsus adductus from other adductory conditions
- 3) To convey the importance of making a proper diagnosis relative to patient management
- 4) To review the pathoanatomy, clinical features, and clinical course of metatarsus adductus
- 5) To review assessment and treatment parameters
- 6) To review, compare, and contrast other adductory conditions that are important in the assessment and management of metatarsus adductus

139

Welcome to Podiatry Management's CME Instructional program. Our journal has been approved as a sponsor of Continuing Medical Education by the Council on Podiatric Medical Education.

You may enroll: 1) on a per issue basis (at \$23.00 per topic) or 2) per year, for the special rate of \$179 (you save \$51). You may submit the answer sheet, along with the other information requested, via mail, fax, or phone. You can also take this and other exams on the Internet at [www.podiatrym.com/cme](http://www.podiatrym.com/cme).

If you correctly answer seventy (70%) of the questions correctly, you will receive a certificate attesting to your earned credits. You will also receive a record of any incorrectly answered questions. If you score less than 70%, you can retake the test at no additional cost. A list of states currently honoring CPME approved credits is listed on pg. 148. Other than those entities currently accepting CPME-approved credit, Podiatry Management cannot guarantee that these CME credits will be acceptable by any state licensing agency, hospital, managed care organization or other entity. PM will, however, use its best efforts to ensure the widest acceptance of this program possible.

**This instructional CME program is designed to supplement, NOT replace, existing CME seminars.** The goal of this program is to advance the knowledge of practicing podiatrists. We will endeavor to publish high quality manuscripts by noted authors and researchers. If you have any questions or comments about this program, you can write or call us at: **Podiatry Management, P.O. Box 490, East Islip, NY 11730, (631) 563-1604 or e-mail us at [bblock@podiatrym.com](mailto:bblock@podiatrym.com).**

Following this article, an answer sheet and full set of instructions are provided (pg. 148).—Editor

**T**he most common inquiries from practitioners concern whether metatarsus adductus in a particular patient requires therapeutic intervention at all, whether

the patient is age-appropriate for treatment, whether there are associated torsional or rotational conditions, or if detected, whether or not this condition even needs to be addressed. Fortunately, current technology

enables remote viewing, so that gait video (where age appropriate), clinical photos, and radiographs are available for assessment. Unfortunately, too often the condition has

*Continued on page 140*

*Differential (from page 139)*

been misdiagnosed.

Practitioners who are not readily familiar with pediatric norms at given stages of development sometimes miss subtle but tell-tale findings that are important in assessment and treatment. This article will focus on metatarsus adductus as well as the differential diagnosis of adductory conditions that may have a component of metatarsus (or forefoot) adductus. There are, however, different conditions with different clinical courses which require different interventions. The reader is reminded that much of the information on the specific conditions referenced is readily available through the Internet and s/he is encouraged to review that information.

### Metatarsus Adductus

Metatarsus adductus is the most common congenital foot condition encountered. The foot is “C-shaped”



Figure 3 (left): Clinical photo of foot in relaxed position, with adducted forefoot.

Figure 4 (right): Clinical photo of forefoot manipulated into rectus position on mid- and hindfoot. Note that abduction is performed at the metatarsal head level rather than at the toes, while the hindfoot is maintained in neutral position.

dren. Note that in a small percentage of cases, the condition may be associated with hip dysplasia, and the practitioner should include clinical assessment of hip stability (e.g. Barlow's, Ortolani's, Galeazzi tests) as part of the overall work-up.

### Clinical Presentation

The clinical presentation is uniplanar, with adduction of the forefoot on the mid- and hindfoot.



Figure 5: Clinical photo of child in stance, demonstrating bilateral metatarsus adductus with concurrent hallux adductus, a common finding.

## In infants, metatarsus adductus and forefoot adductus are indistinguishable because the lesser tarsus is radiographically silent.

with a convex lateral border and concave medial border (Figure 1). It is most likely caused by intrauterine position (intrauterine packing); however, the literature reports some association with oligohydramnios (lack of amniotic fluid). It is present at birth and most often noted through the first year. The vast majority of cases are flexible (90%); however, a spectrum of flexibility exists, and semi-rigid as well as frankly rigid conditions are encountered.

Approximately 50% of presentations are bilateral, though not necessarily perfectly symmetric. Flexible presentations typically resolve spontaneously in approximately 85% of chil-

While accurately represented as adduction of the metatarsals on the lesser tarsus, it is clinically indistinguishable



Figure 1: Clinical photo of metatarsus adductus (from plantar view), revealing classic findings of concave medial border and convex lateral border, with the apex of the deformity at the base of the fifth metatarsal.



Figure 2: AP radiograph of metatarsus adductus, demonstrating age-appropriate divergence of Kite's angle, radiographically “silent” lesser tarsus, and adduction of metatarsals. The talar bisection projects lateral to the bisection of the first metatarsal.

from forefoot adductus in pre-walkers and beyond. The midfoot is radiographically “silent”, meaning that the ossific landmarks of the cuboid and cuneiforms are either not present or not yet well visualized (Figure 2). The ossific nuclei of the tarsals, however, are present.

As a result, the positional relationship between the metatarsals and lesser tarsals cannot be accurately determined. In addition, with the vast majority of cases being flexible, observation and management are undertaken without the need for radiographs. Practically speaking, the preponderance of children presented to the practitioner fall into the pre-walker age bracket, all of which share the feature of radiographic “silence”. Therefore, in pre-walkers it falls primarily to the salient assessment of clinical features, including cutaneous landmarks, ranges of motion of the fore-

*Continued on page 141*




Figure 7 (above): Lateral radiograph demonstrating typical angular relationships in metatarsus adductus, where the lesser tarsus can be visualized.

Figure 6 (left): AP radiograph demonstrating typical angular relationships in metatarsus adductus, where the lesser tarsus can be visualized.

Where the condition persists to the point that radiographic landmarks are visualized, the following values are typical:

AP/DP view (Figure 6):

- Kite's angle WNL
- talar-1st metatarsal angle  $> 15^\circ$  (N =  $< 15^\circ$ )
- calcaneo-1st metatarsal angle  $> 10^\circ$  (N =  $0 \sim 5^\circ$ )
- calcaneo-2nd metatarsal angle  $> 10^\circ$  (N =  $0 \sim 5^\circ$ )

Lateral view (Figure 7):

- calcaneal inclination angle: WNL (N =  $18 \sim 25^\circ$ )
- talar declination angle: WNL (N =  $15^\circ - 25^\circ$ )
- talar-1st metatarsal angle: WNL (N =  $15^\circ - 25^\circ$ )

### Differential Diagnosis

While it is important to assess the flexibility of metatarsus adductus relative to possible intervention, it is

### Differential (from page 140)

foot, subtalar and ankle joints, as well as transverse rotation of the leg on the thigh, and hip range of motion, to determine whether additional diagnoses should be considered, as well as the need for radiographs.

A useful clinical parameter in foot assessment is the ease of manual reduction of the forefoot on the hindfoot. Flexible presentations are typified by the forefoot being easily manipulated into a rectus or mildly abducted position (Figures 3 and 4).

Care should be taken to abduct rather than pronate the forefoot, as pronation of the forefoot will lead to a "spurious" reduction and mislead the practitioner into a false sense of its responsiveness to manipulation. This is even more important in semi-rigid or rigid conditions. In the latter two, the practitioner should evaluate cutaneous landmarks, such as creases in the mid-arch and poste-

is undertaken, a short-leg cast is as effective as a long-leg cast.

Use of long-leg casts for isolated metatarsus adductus is more pragmatic than therapeutic, especially when dealing with a chubby infant or an older child whose repetitive knee motion may shift the position of the

**In a six month old, the most common way to differentiate metatarsus adductus from skew foot is the extent of subtalar pronation.**

cast on the leg. Where concurrent internal tibial torsion is present (as it frequently is), the practitioner may wish to employ a long-leg cast to correct the conditions simultaneously. Note, however, that as with metatarsus adductus, mild super-structural rotational disorders typically resolve spontaneously.

equally important to understand other "adductory" conditions that cosmetically mirror it but that have more significant consequence if missed by the practitioner. They are talipes equinovarus (TEV or clubfoot), skew foot, and to a lesser extent internal tibial torsion, and femoral anteversion.

### Talipes Equinovarus

Talipes equinovarus (TEV, or clubfoot) shares forefoot adduction, clinically, with metatarsus adductus; however, like skew foot, it is a complex deformity.

The condition is typified by:

- Adduction of the forefoot and midfoot on the hindfoot (not metatarsal adduction on the lesser tarsus). A common distinguishing characteristic in TEV, especially in more severe cases, is a medial fissure (crease) in the mid-arch area (Figure 8), and often a fissure (crease) at the posterior ankle (Figures 9 and 10). Radio-

*Continued on page 142*

### The typical clinical course

**for metatarsus adductus resolves spontaneously in most cases.**

rior ankle, as these are common with talipes equinovarus. If this is ruled out, intervention via manipulation with serial casting may be required, as over time articular positions and soft tissue adaptation become less amenable to spontaneous reduction with weight-bearing. If serial casting

In older children who are fully weight-bearing and ambulating, pronatory compensation may cause mild FF supinatus with mild hindfoot valgus (important to distinguish from skewfoot). The hallux is often adducted as well and can be exaggerated with weight-bearing (Figure 5).



*Differential (from page 141)*

graphically, however, both conditions will demonstrate bisection of the first metatarsal falling medial to a bisection of the talus on an AP view (Figures 11 and 12).

- Supination of the subtalar joint. Clinically, the hindfoot will show little range of motion in the frontal plane. Metatarsus adductus demonstrates typical pediatric values (mild increase) for Kite's angle. Tarsal stacking is pathognomonic of TEV and will readily differentiate it radiographically from metatarsus adductus (Figure 11). In addition, a lateral radiograph of TEV (Figure 12) will demonstrate lack of anterior convergence of the talar and calcaneal bisections. The talar bisection will fall in the dorsal half of the ossific nucleus of the cuboid and often project entirely dorsal to it. In more

subtalar capsule, the calcaneofibular ligament and the posterior talofibular ligament. Concurrently, there is gastrocnemius contracture. This combination of contractures maintains the calcaneal pitch low or even negative.

us should be manually reduced via the Ponseti method, which entails a different series of maneuvers and hand positions. The reader is referred to the literature and Internet to review the Ponseti method of

### **In skewfoot, Kite's angle is expected to be greater than that of metatarsus adductus.**

In contrast, with metatarsus adductus, the hindfoot equinus is atypical, and when present, is much more mild and commonly due to gastrocnemius contracture.

The functional significance of these differences is important in management. If a true TEV is misdiagnosed as metatarsus adductus and manipulation with serial casting is

manipulation, as it is beyond the scope of this article.

If surgical correction is undertaken for what is misdiagnosed as a metatarsus adductus but is truly a TEV, the practitioner will fail to reduce the hindfoot varus, fail to perform a tendo Achilles lengthening and, if required, fail to release the posterior capsules and ligaments. As a result, the forefoot adductus correction will not be maintained, and the hindfoot will remain in equinovarus. Even worse, if a tendo Achilles lengthening is performed but the hindfoot varus has not been addressed, an iatrogenic cavaoadducto-varus is likely.

### **The presence of a posterior calcaneal fissure is indicative of hindfoot equinus.**

severe TEV, parallelism of the bisections of the talus and calcaneus is present (Figure 12).

- Hindfoot equinus is another key characteristic that differentiates TEV from metatarsus adductus. In TEV, it is invariably present, because of the posterior calcaneus being drawn upward due to contracture of the posterior ankle joint capsule, the posterior

undertaken, the practitioner, in an attempt to abduct the metatarsals on the lesser tarsus, will encounter resistance. The undetected hindfoot varus will cause resistance to the abduction. This is because the forefoot adduction in TEV is occurring at Chopart's joint and not Lisfranc's joint.

Unfortunately, if the practitioner is persistent, the tendency is to subluxate the lateral column. A true talipes equinovar-

#### **Skewfoot**

Skewfoot, also referred to as "Z foot", is typified by three areas of deviation:

- 1) The forefoot is adducted on the midfoot, as with metatarsus adductus. Depending on severity, the first metatarsal lies
- 2) adducted and
- 3) plantar-flexed on the medial cuneiform.

- In contrast to metatarsus adductus, the midfoot is abducted relative to the hindfoot, with lateral displacement of the navicular on the head of the talus and abduction of the cuboid on the calcaneus (Figure 11). In more severe cases, the navicular may also lie dorsally subluxed in relation to the talus.

Note that in the radiograph shown, the ossific nucleus of the cuboid is

*Continued on page 143*



Figure 8: Clinical photo of talipes equinovarus (from plantar view), demonstrating forefoot adduction. Note the skin crease in the medial arch.



Figure 9: Clinical photo of talipes equinovarus (from anterior view), demonstrating hindfoot varus (hindfoot supination).



Figure 10: Clinical photo of talipes equinovarus (from medial view), demonstrating skin crease at superior heel area, consistent with hindfoot equinus.



Figure 11: AP radiograph of talipes equinovarus, demonstrating superimposition of the talus on the calcaneus (tarsal stacking), and indicative of hindfoot supination.

Figure 12 (right): Lateral radiograph of talipes equinovarus, demonstrating superimposition of the talus on the calcaneus (tarsal stacking), and indicative of hindfoot supination. Note parallelism of talus and calcaneus, with rotation of the calcaneus into equinus.

[Clinical Pearl: Note that to accommodate positioning of the lateral border of the foot against the plate, the hindfoot becomes relatively adducted and the ankle becomes more externally rotated. The ankle joint therefore appears as a mortise view while the talus and calcaneus appear foreshortened. Alternatively, the foot can be positioned with the lateral hindfoot against the plate, which will render a true representation of the size and relationship of the tarsal bones, as well as a “true” projection across the ankle. The trade-off is that the forefoot will become distorted.



cally “rectus” forefoot, but a significantly pronated hindfoot.

If a misdiagnosis of metatarsus adductus is made in the case of a resistant or rigid deformity and surgical correction is undertaken, the practitioner will correct the forefoot and again neglect the abducted midfoot and pronated hindfoot. The result will be a partial correction which will destabilize the foot medially. Weight-bearing children with skewfoot rely on the forefoot adduction for medial stability in stance. Failure to address all components of the deformity concurrently and focus only on the forefoot adduction will

exacerbate midfoot abduction and hindfoot pronation when the patient is weight-bearing.

Comparative AP radiographs of metatarsus adductus, TEV and skewfoot are shown (Figure 15). Note that while the metatarsals are adducted in all three conditions, the osseous positions at the lesser tarsus and tarsus are significantly different among the three.

#### Internal Tibial Torsion

Internal tibial torsion is a normal condition in infants and is present through 18-24 months of age. It is defined by the relationship of the leg to the thigh and typically resolves spontaneously. It is easily assessed by internally and externally rotating the leg on the thigh with the knee flexed

### When serial casting is employed in a patient in whom a skewfoot is undetected (misdiagnosed as metatarsus adductus), the hindfoot will most likely be iatrogenically supinated in the cast.

#### Differential (from page 142)

present but that of the navicular does not yet show. The diagnostic challenge occurs in more mild cases of skewfoot in infants, pre-walkers and early walkers because the ossific nuclei of the navicular and cuboid are either not visualized or, if visualized, are not well enough delineated radiographically to determine their abductory position relative to the tarsal bones (which are also poorly defined radiographically at this age).

- In contrast to metatarsus adductus, the hindfoot is in valgus, with pronation and concurrent adduction and plantarflexion of the talus. In more severe cases, calcaneal pitch is low, with tight gastrocnemius or gastrosoleus (i.e., tendo Achilles contracture). In pre-walkers, however, the positional valgus may not be readily apparent and the practitioner should carefully evaluate the subtalar range of motion.

The functional significance of these differences is important in man-

agement. If a misdiagnosis of metatarsus adductus is made, and manipulation with serial casting is undertaken, the practitioner, in the attempt to abduct the metatarsals on the lesser tarsus, will instead pronate the hindfoot, as the Lisfranc articulations are the most resistant to correction. Therefore, care must be taken to prevent hindfoot pronation with counter pres-

### When serial casting is employed in patient in whom a TEV is undetected (misdiagnosed as metatarsus adductus), the metatarsals will resist long-term correction due to hindfoot supination.

sure at the hindfoot, and to further maintain this in the cast. Otherwise, the subtalar joint, already over-pronated in skewfoot, will continue to pronate, and the practitioner risks a spurious correction in the transverse plane, with the end result of a clini-

at 90° and the foot held in neutral. Another commonly used static assessment is the thigh-foot angle (Figure 16). This angle relates to a bisection of the posterior thigh to the centerline of the foot.

*Continued on page 144*

*Differential (from page 143)*

Where the thigh-foot angle is internal and exceeds -10 degrees at 24 months, monitoring (though not necessarily treatment) is warranted. Inter-

the foot level only, are less effective and tend to shift the moment of force to the femur rather

## Internal tibial torsion is a normal condition in infants and is present through 18-24 months of age.

vention, where indicated, is most often accomplished with a Wheaton-type brace, which appropriately maintains an abductory force on the tibia with the knee in a flexed position. Older therapies, such as abductory bars at




Figure 13: AP radiograph of skewfoot, demonstrating increased Kite's angle, with abduction of the cuboid on the calcaneus, as well as adduction of the metatarsals.




Figure 14: Clinical photo of skewfoot, demonstrating forefoot findings similar in appearance to metatarsus adductus, but significant fibular displacement of the lesser tarsus on relative to the leg, concurrent with hindfoot pronation.

er than where it is intended.

The functional significance in a discussion of metatarsus adductus is that the two conditions often present in tandem. Flexible conditions are of less concern, since both resolve spontaneously in the vast majority of cases. Where either or both persist beyond their natural course, the practitioner needs to be able to anticipate the presence of both, and address management specifically to the areas where it will be effective. As previously indicated, serial long-leg casting will incorporate the appropriate positional abduction at the foot level,

presentation of the child. The condition's normal course results in near complete resolution of the in-toeing in more than 80% of affected children by age 10.

Concurrent metatarsus adductus and femoral anteversion is less common than is metatarsus adductus with internal tibial torsion. The functional significance for the practitioner is in making a thorough segmental assessment when faced with the tripping child, or with the child who is presented for "in-toeing." A simple assessment method in ambulatory




Figure 15: Comparative AP radiographs of metatarsus adductus, talipes equinovarus, and skewfoot, outlining the salient tarsal relationships and position relative to the cuboid and forefoot.

## A simple assessment method in ambulatory children to detect femoral anteversion is to observe the patellae as the child approaches during gait.

while also maintaining an abductory position at the tibia. When followed by bracing (e.g., Whitman brace), the two areas will be incorporated in therapy.

### Femoral Anteversion

Femoral anteversion (medial femoral torsion) is the most common cause of in-toeing in early childhood (Figure 18).

It is defined by the relation between the femoral neck and the transcondylar axis at the knee. It is normal at birth and reduces with age, but if delayed may be a cause of tripping or awkward gait in children, and is commonly the reason for parental

children is to observe the patellae as the child approaches during gait. The patellae will be internal to the sagittal plane.

Femoral anteversion may be the source, or a contributory element, of the child's adducted angle of gait. The practitioner should be aware that children commonly pronate in compensation for the adduction, and in so doing, may be masking a persistent metatarsus adductus. The practitioner should therefore assess the feet in their compensated position as well as with the child placed in corrected stance position, with both feet placed in neutral stance. If a metatarsus ad-

*Continued on page 145*



*Differential (from page 144)*

ductus (or forefoot adductus) is present, it will become more evident.

### Treatment Summary for Metatarsus Adductus

#### Non-Operative

- Defer treatment if the foot can be passively manipulated past neutral (rectus lateral border).
- Some patients may benefit from passive stretching.
- Adjunctive therapies include bar therapy, straight last or abductor shoes, pre-fabricated AFOs, etc.; while broadly utilized in the past, strong medical evidence is lacking for these interventions.

rection is obtained, several holding casts are advised to maintain correction.

#### Operative

- Abductor hallucis release, capsulotomy, and metatarsal osteotomy are surgical options for young children; adjunctive cuboid osteotomy has been effectively performed as well.
- Tarsometatarsal capsulotomies are still utilized; however, they should be relegated to young children, as stiffness is common in children five years or older.
- In patients older than five years




Figure 16: Illustration of the thigh-foot axis.

Orthop. 1983 Feb;3(1):2-9.

7) Calcaneal lengthening for valgus deformity of the hindfoot. Results in children who had severe, symptomatic flatfoot and skewfoot.; Mosca VS., J Bone Joint Surg Am. 1995 Apr;77(4):500-12.

8) Resistant metatarsus adductus: prospective randomized trial of casting versus orthosis., Herzenberg JE, Burghardt, RD, J Orthop Sci. 2014 Mar;19(2):250-6.

Metatarsus adductus: development of a non-surgical treatment pathway. Williams CM, James AM, Tran T., J Paediatr Child Health. 2013 Sep;49(9):E428-33.

9) A reappraisal of metatarsus adductus and Skewfoot, Berg EE. J Bone Joint Surg Am. 1986 Oct; 68(8):1185-96.

10) Skewfoot; Foot Ankle Clin. 2009 Hagmann S, Dreher T, Wenz W. Sep;14(3):409-34.

12) Results of clubfoot management using the Ponseti method: do the details matter? A systematic review.; Zhao D, Li H, Zhao L, Liu J, Wu Z, Jin F.; Clin Orthop Relat Res. 2014 Apr;472(4):1329-36.

13) Tibial torsion: a method of assessment and a survey of normal children. Staheli LT, Engel GM., Clin Orthop Relat Res. 1972 Jul-Aug;86:183-6.

14) Common rotational variations in children. Lincoln TL, Suen PW. J Am Acad Orthop Surg. 2003 Sep-Oct; 11(5):312-20.

15) The normal development of tibial torsion. Kristiansen LP, Gunderson RB, Steen H, Reikerås O.; Skeletal Radiol. 2001 Sep;30(9):519-22.

16) Measurement of tibial torsion and thigh-foot angle using goniometry and computed tomography. Stueberg W, Temme J, Kaplan P, Clarke A, Fuchs R. Clin Orthop Relat Res. 1991 Nov; (272):208-12.

## A Wheaton brace is indicated in a one year-old with resistant metatarsus adductus and internal tibial torsion.

• Serial casting is indicated for rigid or persistent deformity; intervention is best started before the child reaches one year.

• Persistent or rigid forefoot adductus can be readily corrected w/ cast; avoid pronating the forefoot with abduction.

• When concurrent internal tibial torsion is present, casts should extend above the knee (flexed 20-25°); otherwise below-knee casting is adequate.

• Casts are changed biweekly, and correction is usually achieved after two to three changes; once cor-

rection is obtained, consider proximal osteotomies of the lesser metatarsals with or without cuboid osteotomy. **PM**

#### Suggested Reading

1) The intoeing child: etiology, prognosis, and current treatment options. Harris E. Clin Podiatr Med Surg. 2013 Oct;30(4):531-65.

2) The long-term functional and radiographic outcomes of untreated and non-operatively treated metatarsus adductus. Farsetti, P, Weinstein SL, Ponseti IV; J Bone Joint Surg Am. 1994 Feb;76(2):257-65.

3) Below-knee plaster cast for the treatment of metatarsus adductus. Katz K, David R, Soudry M. J Pediatr Orthop. 1999 Jan-Feb; 19(1):49-50.

4) Radiological assessment of metatarsus adductus. Dawoodi AI, Perera A. Foot Ankle Surg. 2012 Mar; 18(1):1-8. Epub 2011 Apr 8.

5) A comparison of two non-operative methods of idiopathic clubfoot correction: the Ponseti method and the French functional (physiotherapy) method. Surgical technique.; Steinman S, Richards BS, Faulks S, Kaipus K.; J Bone Joint Surg Am. 2009 Oct 1;91 Suppl 2:299-312.

6) Metatarsus adductus: classification and relationship to outcomes of treatment; Bleck EE. J Pediatr



Figure 17: Clinical photo of a child with internal tibial torsion. Note the position of the foot relative to the patella.



Figure 18: Clinical photo of a child with internal femoral position. Note the adducted position of the patellae and the foot angle in stance.



**Dr. Benard** is Executive Director of the American Board of Podiatric Medicine and Co-Director of the Baja Project for Crippled Children. He is a Diplomate of the American Board of Podiatric Medicine, and

the American Board of Podiatric Surgery and lectures nationally on topics in biomechanics and pediatric foot and ankle surgery. He is also the current Assistant-Governor for International Service and Disaster Relief for Rotary International District 5280.

## SEE ANSWER SHEET ON PAGE 149.

- 1) Approximately what percent of metatarsus adductus cases are bilateral?
  - A) 5%
  - B) 50%
  - C) 90%
  - D) 99%
- 2) In infants, metatarsus adductus and forefoot adductus are indistinguishable because:
  - A) The lesser tarsus is radiographically silent
  - B) The tarsal position is the same in either condition
  - C) Both are uniplanar
  - D) The cuboid is abducted on the calcaneus in both conditions
- 3) The typical clinical course for metatarsus adductus is:
  - A) Persists and causes compensation at the subtalar joint
  - B) Resolves spontaneously in most cases
  - C) Requires a course of serial casting
  - D) Prognosis is poor when concurrent internal tibial torsion is present
- 4) In a six-month-old, the most common way to differentiate metatarsus adductus from skew foot is:
  - A) Rigidity of forefoot adduction
  - B) Presence of gastrosoleus contracture
  - C) Talus is superimposed on the calcaneus
  - D) Extent of subtalar pronation
- 5) The presence of a posterior calcaneal fissure is indicative of:
  - A) Limitation of subtalar range of motion
  - B) Intrauterine position
  - C) Hindfoot equinus
  - D) Rigid metatarsus adductus
- 6) On reviewing Figures 3 and 4, which of the following constitutes appropriate management?
  - A) Serial casting should be initiated
  - B) Employ Ponseti technique
  - C) Defer treatment and monitor
  - D) Supinate hindfoot in cast
- 7) In skewfoot, Kite's angle is expected to be:
  - A) Less than that of metatarsus adductus
  - B) Less than that of talipes equinovarus
  - C) Greater than that of metatarsus adductus
  - D) Equal to that of talipes equinovarus
- 8) When serial casting is employed in a patient in whom a skewfoot is undetected (misdiagnosed as metatarsus adductus):
  - A) The hindfoot will most likely be iatrogenically supinated in the cast
  - B) The hindfoot will most likely be iatrogenically pronated
  - C) The metatarsals will be the most responsive to correction
  - D) The hindfoot will iatrogenically be plantarflexed
- 9) When serial casting is employed in a patient in whom a TEV is undetected (misdiagnosed as metatarsus adductus):
  - A) The hindfoot will most likely be iatrogenically supinated in the cast
  - B) Iatrogenic abduction of the cuboid may occur
  - C) The hindfoot will be iatrogenically plantarflexed
  - D) The metatarsals will resist long-term correction due to hindfoot supination
- 10) A simple assessment method in ambulatory children to detect femoral anteversion is to:
  - A) Observe the patellae as the child approaches during gait
  - B) Observe for cadence differential in unilateral cases
  - C) Observe for pelvic tilt
  - D) Observe for hyperlordosis
- 11) Which of the following is an appropriate method of assessing tibial torsion?
  - A) Evaluating the thigh-foot axis
  - B) Noting the response to serial casting after two applications
  - C) Employing Barlow's examination
  - D) None of the above
- 12) Which of the following radiographic values are inconsistent with metatarsus adductus?
 

AP/DP view:

  - A) Kite's angle WNL
  - B) Calcaneo-1st metatarsal angle greater than 10°
  - C) Calcaneo-2nd metatarsal angle greater than 10°
  - D) All of the above
- 13) Which of the following radiographic values are consistent with metatarsus adductus?
 

Lateral view:

  - A) Talar declination angle: WNL
  - B) Talar-1st metatarsal angle WNL
  - C) Choices A & B
  - D) None of the above
- 14) A seven-month-old is presented for "intoeing." A course of serial casting was performed by another practitioner at age one month. Four cast changes were performed at two week intervals. The mother reports initial correction, but over time the in-toeing recurred. You have ruled out torsional/rotational factors, and have obtained the radiographs shown in Figures 11 and 12.
 

Which of the following is the most likely reason for failure of prior treatment?

  - A) Casting was initiated too early
  - B) More manipulation and

Continued on page 148



cast applications were required

- C) An underlying neurologic condition was present
- D) Undetected hindfoot varus resisted abduction of the forefoot

15) In the patient referenced in Question #14, current management should include:

- A) Reinstitution manipulation via the Ponseti technique
- B) Reinstitution manipulation, and both pronate and abduct the forefoot
- C) Delay further manipulation until the lesser tarsus is better visualized
- D) Delay further manipulation and perform metatarsal osteotomies

16) Internal tibial torsion is a normal condition in infants and is present through \_\_\_\_\_ months of age.

- A) 1-6
- B) 6-12
- C) 12-18
- D) 18-24

17) Serial casting of the foot should, in addition to abduction of the forefoot, include:

- A) Abduction of the midfoot on the hindfoot
- B) Maintaining the hindfoot in neutral
- C) No additional positioning
- D) Plantar-flexion of the ankle

18) Which of the radiographs in Figure 15 is consistent with talipes equinovarus?

- A) Left
- B) Center
- C) Right
- D) None of the above

19) In metatarsus adductus, the foot is:

- A) A-shaped
- B) C-Shaped
- C) O-shaped
- D) S-Shaped

20) Which of the following is most indicated in a one-year-old with resistant metatarsus adductus and internal tibial torsion?

- A) Denis-Browne bar
- B) Straight last shoe
- C) Wheaton brace
- D) Short-leg cast

## PM's CME Program

Welcome to the innovative Continuing Education Program brought to you by *Podiatry Management Magazine*. Our journal has been approved as a sponsor of Continuing Medical Education by the Council on Podiatric Medical Education.

### Now it's even easier and more convenient to enroll in PM's CE program!

You can now enroll at any time during the year and submit eligible exams at any time during your enrollment period.

PM enrollees are entitled to submit ten exams published during their consecutive, twelve-month enrollment period. Your enrollment period begins with the month payment is received. For example, if your payment is received on November 1, 2014, your enrollment is valid through October 31, 2015. If you're not enrolled, you may also submit any exam(s) published in PM magazine within the past twelve months. **CME articles and examination questions from past issues of *Podiatry Management* can be found on the Internet at <http://www.podiatrym.com/cme>.** Each lesson is approved for 1.5 hours continuing education contact hours. Please read the testing, grading and payment instructions to decide which method of participation is best for you.

Please call (631) 563-1604 if you have any questions. A personal operator will be happy to assist you.

Each of the 10 lessons will count as 1.5 credits; thus a maximum of 15 CME credits may be earned during any 12-month period. You may select any 10 in a 24-month period.

**The Podiatry Management Magazine CME program is approved by the Council on Podiatric Education in all states where credits in instructional media are accepted. This article is approved for 1.5 Continuing Education Contact Hours (or 0.15 CEU's) for each examination successfully completed.**

**Home Study CME credits now  
accepted in Pennsylvania**

**SEE ANSWER SHEET ON PAGE 149.**

# Enrollment/Testing Information and Answer Sheet

Continuing  
Medical Education

**Note:** If you are mailing your answer sheet, you must complete all info. on the front and back of this page and mail with your credit card information to: **Podiatry Management, P.O. Box 490, East Islip, NY 11730.**

## TESTING, GRADING AND PAYMENT INSTRUCTIONS

(1) Each participant achieving a passing grade of 70% or higher on any examination will receive an official computer form stating the number of CE credits earned. This form should be safeguarded and may be used as documentation of credits earned.

(2) Participants receiving a failing grade on any exam will be notified and permitted to take one re-examination at no extra cost.

(3) All answers should be recorded on the answer form below. For each question, decide which choice is the best answer, and circle the letter representing your choice.

(4) Complete all other information on the front and back of this page.

(5) Choose one out of the 3 options for testgrading: mail-in, fax, or phone. To select the type of service that best suits your needs, please read the following section, "Test Grading Options".

## TEST GRADING OPTIONS

### Mail-In Grading

To receive your CME certificate, complete all information and mail with your credit card information to:

**Podiatry Management**

**P.O. Box 490, East Islip, NY 11730**

**PLEASE DO NOT SEND WITH SIGNATURE REQUIRED, AS THESE WILL NOT BE ACCEPTED.**

There is **no charge** for the mail-in service if you have already en-

rolled in the annual exam CME program, and we receive this exam during your current enrollment period. If you are not enrolled, please send \$23.00 per exam, or \$179 to cover all 10 exams (thus saving \$51 over the cost of 10 individual exam fees).

### Facsimile Grading

To receive your CME certificate, complete all information and fax 24 hours a day to 1-631-563-1907. Your CME certificate will be dated and mailed within 48 hours. This service is available for \$2.50 per exam if you are currently enrolled in the annual 10-exam CME program (and this exam falls within your enrollment period), and can be charged to your Visa, MasterCard, or American Express.

If you are *not* enrolled in the annual 10-exam CME program, the fee is \$23 per exam.

### Phone-In Grading

You may also complete your exam by using the toll-free service. Call 1-800-232-4422 from 10 a.m. to 5 p.m. EST, Monday through Friday. Your CME certificate will be dated the same day you call and mailed within 48 hours. There is a \$2.50 charge for this service if you are currently enrolled in the annual 10-exam CME program (and this exam falls within your enrollment period), and this fee can be charged to your Visa, Mastercard, American Express, or Discover. If you are not currently enrolled, the fee is \$23 per exam. When you call, please have ready:

1. Program number (Month and Year)
2. The answers to the test
3. Your social security number
4. Credit card information

In the event you require additional CME information, please contact PMS, Inc., at **1-631-563-1604**.

149

## ENROLLMENT FORM & ANSWER SHEET

*Please print clearly...Certificate will be issued from information below.*

Name \_\_\_\_\_ Soc. Sec. # \_\_\_\_\_

Please Print: FIRST MI LAST

Address \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Charge to: ☐ Visa ☐ MasterCard ☐ American Express

Card # \_\_\_\_\_ Exp. Date \_\_\_\_\_

**Note: Credit card is the only method of payment. Checks are no longer accepted.**

Signature \_\_\_\_\_ Soc. Sec. # \_\_\_\_\_ Daytime Phone \_\_\_\_\_

State License(s) \_\_\_\_\_ Is this a new address? Yes ☐ No ☐

**Check one:** ☐ I am currently enrolled. (If faxing or phoning in your answer form please note that \$2.50 will be charged to your credit card.)

☐ I am not enrolled. Enclosed is my credit card information. Please charge my credit card \$23.00 for each exam submitted. (plus \$2.50 for each exam if submitting by fax or phone).

☐ I am not enrolled and I wish to enroll for 10 courses at \$179.00 (thus saving me \$51 over the cost of 10 individual exam fees). I understand there will be an additional fee of \$2.50 for any exam I wish to submit via fax or phone.

**EXAM #7/14**  
**Differential Diagnosis of Metatarsus Adductus**  
**(Benard)**

**Circle:**

- |             |             |
|-------------|-------------|
| 1. A B C D  | 11. A B C D |
| 2. A B C D  | 12. A B C D |
| 3. A B C D  | 13. A B C D |
| 4. A B C D  | 14. A B C D |
| 5. A B C D  | 15. A B C D |
| 6. A B C D  | 16. A B C D |
| 7. A B C D  | 17. A B C D |
| 8. A B C D  | 18. A B C D |
| 9. A B C D  | 19. A B C D |
| 10. A B C D | 20. A B C D |

**Medical Education Lesson Evaluation**

Strongly agree [5]	Agree [4]	Neutral [3]	Disagree [2]	Strongly disagree [1]
--------------------------	--------------	----------------	-----------------	-----------------------------

- 1) This CME lesson was helpful to my practice \_\_\_\_
- 2) The educational objectives were accomplished \_\_\_\_
- 3) I will apply the knowledge I learned from this lesson \_\_\_\_
- 4) I will make changes in my practice behavior based on this lesson \_\_\_\_
- 5) This lesson presented quality information with adequate current references \_\_\_\_
- 6) What overall grade would you assign this lesson?  
A B C D

How long did it take you to complete this lesson?  
\_\_\_\_ hour \_\_\_\_ minutes

What topics would you like to see in future CME lessons ?  
Please list :

---



---



---



---



---