



# Prevention of Lower Extremity Amputations

Conservative modalities can lessen morbidity.

BY PRIYANKA BEGUR, DPM, MA AND ROBERT G. FRYKBERG, DPM, MPH

Amputations have long been feared as one of the most tragic complications of diabetes mellitus. Non-healing diabetic foot ulcers (DFU) are most often key components in the pathway leading to non-traumatic amputation in this population. Although frequently necessary as a curative measure for chronic wounds, infections, and/or non-reconstructable ischemia, modern wound care protocols have been established to aim at minimizing the need for amputation. Once healed, prevention programs must focus on appropriate foot care, footwear, and early diagnosis of impending problems in order to avoid new or recurrent lesions. This article briefly outlines modern approaches to the management of DFUs and a new technology to aid in early detection.

Lower extremity amputations are one of the oldest surgical procedures that were performed, dating back to prehistoric times.<sup>1</sup> Greater than one million people with diabetes suffer a limb loss, which equates to an amputation occurring every 20 seconds.<sup>2</sup> Amputations may be indicated for limbs with peripheral vascular disease, trauma, tumors, infections, and congenital limb deficiencies. Peripheral vascular disease is the leading indication for lower extremity amputation in the United States.<sup>3</sup> This is often seen in individuals with diabetes mellitus and peripheral neuropathy. Approximately 65,000 lower extremity amputations are a result of ischemic disease, leading to ulcerations and

subsequent gangrene.<sup>4</sup> It was also reported that 84% of lower extremity amputations are preceded by an ulceration.<sup>5</sup> Though amputations in various cases are considered a limb salvaging technique and can be the best option for a compromised limb, it is vital to treat patients with all appropriate conservative methods prior to resorting to this surgical option.

Approximately 60% of both diabetic and non-diabetic, non-traumatic amputations that are performed might have been prevented with appropriate

tation prevention programs including early pressure detection and annual check-ups in patients at high risk for ulcerations.

## Basic Tenets of Wound Care

Identifying ulcerations, as well as continuing wound care of already identified ulcerations are vital components to preventing amputations. Routine foot checks are one way to help identify and follow ulcers. Patients who are at high risk for developing ulcerations should have rou-

---

**Non-healing diabetic foot ulcers (DFU) are most often key components in the pathway leading to non-traumatic amputation in this population.**

---

and timely care. It has been reported that three key factors can help reduce the rate of amputation.<sup>6</sup> These include appropriate patient education, routine clinic visits with comprehensive foot care, and appropriate footwear. With these three factors, it is estimated that lower extremity amputations should reduce by up to 80%. Various conservative in-clinic options aside from routine foot checks, footwear, and patient education, are available prior to resorting to surgical intervention. This article will focus on the basic tenets of wound care, collagen and growth factor-based creams and gels, antimicrobial biofilm-disrupting dressings, topical oxygen treatment, biologic grafting options, and ampu-

tation foot checks throughout the year or at least annually. These especially include patients with co-morbidities such as diabetes, neuropathy, vascular disease, and those who have a history of a prior ulceration. Patients who are avid smokers should also be monitored closely, since increased tobacco use has been noted to decrease tissue oxygenation, lead to inadequate microbial eradication, poor collagen production, and detrimental effects on leukocytes and fibroblasts affecting immune responses.<sup>7</sup>

Routine foot checks should always include a thorough discussion regarding a history of ulcerations, surgeries, and smoking status, as

*Continued on page 48*



## *Amputations (from page 47)*

well as other co-morbidities that can increase the risk of ulcerations. For routine visits, a dermatological inspection of both feet for ulcerations or pre-ulcerations, areas of erythema, signs of infection including malodor and drainage, and areas of hyperker-

recommended to have ABI measurements performed once they reach the age of 50. Those patients with abnormal vascular examinations, a history of DFUs, prior intervention for peripheral vascular disease or atherosclerotic cardiovascular disease should have annual vascular exams performed. Revascularization is rec-

uct and to facilitate proper healing.

With DFUs that fail to demonstrate improvement, defined as > 50% of reduction in the size of the wound after at least four weeks, other wound care therapies should either be added to the current treatment plan as adjunct therapy, or the treatment should be changed altogether. These can include options such as negative pressure therapy, biologics, and oxygen therapy.

---

---

## **Patients with diabetes are recommended to have ABI measurements performed once they reach the age of 50.**

---

---

### **Topical Gels, Creams, Dressings, and Other Biologics**

In routine clinic visits, lower extremity wounds are typically treated with some sort of topical gel or cream and an appropriate dressing, whether it is the only treatment method, or in conjunction with other therapies. It is vital to identify the cause of the wound so that an appropriate treatment option can be employed. However, using appropriate topical wound care products and dressings can help to minimize or eliminate infections and can help prevent future infection.

It is well accepted that the use of silver is a tried and true method in the prevention and treatment of infection. It supplies a broad spectrum of coverage while having very few side effects. Silver ions react by binding to negatively charged particles, including DNA, RNA, and chloride ions, prevent-

atotic tissue should be conducted. A neurological examination determining gross and protective sensation status should always be completed as well. This includes utilizing a monofilament to test for protective sensation, using a tuning fork for vibration testing, and checking for ankle reflexes. A thorough vascular examination utilizing Doppler ultrasound to check for pedal pulses, observing for edema and checking for symptoms of claudication and rest pain should always be conducted. A musculoskeletal examination checking for bony prominences and a biomechanical examination that could be prompting the formation of an ulcer is vital to prevent future ulcers from forming.<sup>8</sup>

In patients with wounds requiring follow-up care, it is recommended that the visits are spaced between one and four-week intervals to monitor reduction of wound size and the status of the healing progress. When infection becomes a concern, it is recommended that initial sharp debridement, antibiotic therapy, and possible surgical intervention is appropriately followed through and in a timely fashion. It is important to follow the guidelines published by the Infectious Diseases Society of America (IDSA) when determining severity of infection and appropriate courses of action.<sup>9</sup>

Peripheral arterial disease (PAD) should be taken into consideration, and patients with diagnosed or suspected PAD should have ankle-brachial index (ABI) measurements performed. Patients with diabetes are

ommended in those individuals with DFUs and PAD.<sup>10</sup>

Off-loading is a mainstay of therapy for wounds in high pressure areas, especially those on the plantar aspects of feet. In patients with plantar diabetic foot ulcerations (DFU), it is recommended to do so with a total contact cast (TCC) or an irremovable ankle walking boot. In patients who require frequent dressing changes, a removable option such as a walking CAM boot is preferred over a TCC. In patients whose wounds are not along the plantar foot, any modality that relieves the pressure at the site of the ulceration is recommended. Patients who are deemed to be at high-risk due to co-morbidities or a history

---

---

## **It is well accepted that the use of silver is a tried and true method in the prevention and treatment of infection.**

---

---

of ulcerations are recommended to continue to wear diabetic footgear or any therapeutic footwear that is pressure-relieving.

In addition to off-loading, various wound care therapies are available to clinicians. Wound care products that help to maintain a moist wound bed while avoiding maceration are recommended. It is recommended that sharp debridement of all devitalized tissue is done first, prior to application of any wound care product to allow for both better absorption of the prod-

ing replication and altering cell membrane permeability.<sup>11</sup> In this way, silver can act as an antimicrobial agent. Both creams and wound dressings containing silver have shown to be effective in the treatment of a wide range of infections, including bacteria, viruses, fungi, and molds. Nonetheless, the use of silver-containing products should be of a limited nature to avoid the development of resistant bacterial species.

Other gels commonly used in clinical settings work by primarily

*Continued on page 49*



## *Amputations (from page 48)*

disrupting the extracellular polymeric substance (EPS) matrix, which is the main component of biofilm. Biofilm is the EPS matrix surrounding microbes, allowing them colony defenses against host immunity, host healing, and most wound care treatments. Biofilm is the “slime layer” that creates a physical barrier limiting chemicals from entering the bacteria. It has been seen that some wound care gels that work by disrupting this biofilm can reduce the number of bacteria present in the wound by 7 to 8 log. A prior study showed that applying a wound gel that disrupts the biofilm for four weeks helped reduce the volume of microbes 32% greater than when treated solely with standard of care. It was also reported that the use of wound gel improved the rate of healing in chronic wounds as 50% greater than with standard of care.<sup>12</sup>

There are also gels that contain recombinant human platelet-derived growth factor. Becaplermin, for example, is produced by recombinant DNA by inserting the gene for B-chains

and is used by clinicians to be aware of various wound care products and their mechanism of action to appropriately decide on a wound care therapy based on the wound and the patient.<sup>13</sup>

---

---

## **It has been reported and widely accepted that topical oxygen for non-healing wounds can have beneficial effects.**

---

---

in platelet-derived growth factors (PDGF). These growth factors induce fibroblast proliferation and differentiation, potentiating healing, especially in those patients with decreased healing capacity. Gels that contain such growth factors that are utilized in ulcerations in conjunction with appropriate wound care, have been seen to heal approximately six weeks faster than those with only standard of care. Therefore, it is important for clini-

### **Topical Oxygen Therapy**

It has been reported and widely accepted that topical oxygen for non-healing wounds can have beneficial effects. The partial pressure of oxygen (pO<sub>2</sub>) is a key factor in wound healing.<sup>14</sup> A pO<sub>2</sub> less than 40 mmHg is associated with poor wound healing. Many chronic non-healing wounds are stuck in the inflammatory phase, due to a lack

*Continued on page 50*



## Amputations (from page 49)

of oxidative killing by neutrophils, which can occur due to a low pO<sub>2</sub> level.<sup>15</sup> Increasing the pO<sub>2</sub> level at the site of a wound helps increase reactive oxygen species, which is a vital component in normal cell signal transduction and cell cycling. In addition, it is seen that adequate oxygen supply regulates cellular motility, angiogenesis, and extracellular matrix formation, all of which are vital for proper wound healing.<sup>16</sup>

Topical oxygen can be administered through various methods. One common method is delivering 100% oxygen for a total of 90 minutes once a day at the site of the wound. Another common method is applying continuous, atmospheric oxygen for three to seven days to the site of the wound through a cannula. Experimental studies have shown that patients using topical oxygen therapy for up to eight days had significantly more epithelialization on histologic examination compared to control wounds that did not have topical oxygen therapy. Various studies on topical oxygen treatment reported that subjects who were placed on topical oxygen in comparison to standard of care treatment healed in a shorter amount of time and had a significantly higher wound size reduction. It is, however, important to note that topical oxygen systems generally are able to diffuse through a maximum distance of 50-100 microns; therefore, the amount of biofilm layers and tissue over the wound should be taken into consideration before utilizing this conservative option.<sup>17</sup>

## Skin Substitutes

Various advancements have occurred in both temporary and permanent skin substitutes. Allografts and xenografts can provide a temporary coverage option; however, they can come with the side-effect of rejection. Bioengineered skin substitutes are available to provide both temporary and permanent coverage, with negligible risk of infection or immunologic response. However, they can be very costly. It is important to appropriately utilize these options based upon the patient and the wound.

Human amniotic membranes have been successfully used to treat chronic cutaneous wounds. They are allografts that exhibit low immunogenicity, reduce inflammation, pain, scarring, and accelerate wound healing.<sup>18</sup> Dehydrated human amnion/chorion membranes (dHACM) contain large numbers of pro-angiogenic growth factors that help with neovascularization and healing within chronic wounds. Clinical trials showed that diabetic foot ulcers treated with dHACM healed 77% at four weeks and 92%

to repair wounds and work as a defense against bacteria, fungi, viruses, and parasites. Clinical studies have shown that after approximately five applications of fish-skin grafts, there was a significant reduction in the absolute wound area.<sup>20,21</sup>

## Thermometric Pressure Detection Devices

Thermometric pressure detection mats are a novel product that have recently become available to help detect both new and recurring ulcers before they appear. Wireless




Figure 1: The above photograph shows the type of thermogenic information that would be sent to the clinician regarding increased pressure points along the plantar aspect of the foot, as detected by a thermometric pressure-detection mat. The image on the left shows the thermogram of a patient on day 12, with an increased temperature reading along the right medial forefoot to the hallux. The image on the right was taken on day 22, after the patient was seen by his clinician and offloaded appropriately in response to the increased pressure reported in the thermogram on the left.

at six weeks when compared to standard of care methods.<sup>19</sup> These results in healing rates and the advancements in allografts with decreased immune responses show promising outcomes.

New skin substitutes have been gaining popularity in clinics, especially due to their low immune response. In addition to porcine and bovine skin substitutes, an acellular fish dermal matrix as an alternative tissue source for chronic, non-healing ulcers is gaining popularity. This product is interesting because of the similar structure to human dermis. It also contains omega-3 fatty acids, thus providing a natural structure to the wound bed with additional bioactive lipids. Acellular fish skin also has antimicrobial peptides that help

floor mats with temperature sensors help to detect increased pressure points along the plantar foot, and transmit information to the manufacturer's monitoring center to help prompt clinicians and their staff to see the patient more routinely and alleviate pressure points in a timely manner. These types of devices detect inflammation that often precedes ulcerations by detecting a difference in temperature between the feet of 2.22°C. In a recent study, it was seen that 86% of patients averaged at least three uses per week. For 34 weeks, 132 patients who had a previously healed diabetic foot ulcer were followed, and it was seen that the mat detected 97% of diabetic foot ulcers five weeks before they were

*Continued on page 52*



## Amputations (from page 50)

clinically observed. Considering the high adherence rate by patients and the extremely high detection rate, these types of novel products that

“Amputation Versus Conservative Treatment of a Stage IV Pressure Ulcer, 1999.

<sup>7</sup> Cope, Graham. The effects of smoking on wound healing. *Wounds UK*. June 2014.

<sup>8</sup> Boulton, Andrew J.M. et al. Comprehensive Foot Examination and Risk Assessment. *Diabetes Care*. 2008 Aug.

ized comparative parallel study of amniotic membrane wound graft in the management of diabetic foot ulcers. *Int Wound J* 2013, 10:502-507.

<sup>20</sup> Mostow EN, Haraway GD, Dalsing M, Hodde JP, King D; OASIS Venus Ulcer Study Group. Effectiveness of an extracellular matrix graft (OASIS Wound Matrix) in the treatment of chronic leg ulcers: a randomized clinical trial. *J Vasc Surg*. 2005;41(5):837-843.

<sup>21</sup> Yang, Chun K, et al. Compassionate Clinical Evaluation of a Novel Acellular Fish-skin Graft which contains omega-3 fatty acids for the closure of hard to heal lower extremity chronic ulcers. *Wounds*. 2016. 28(4): 112-118.

<sup>22</sup> Frykberg, Robert G, Gordon, Alexander M, et al. Feasibility and efficacy of a smart mat technology to predict development of diabetic plantar ulcers. *Diabetes Care*. 2017 Apr.

<sup>23</sup> Ertl, Janos P, Panchbhavi Vinod K. Lower-Extremity Amputations 2016 Apr 4.

<sup>24</sup> Elroy, Weledji P, Fokum, Pius. Treatment of the diabetic foot—to amputate or not? *BMC Surg*. 2014; 14:83.

<sup>25</sup> Koob, Thomas J, Lim, Jeremy J, et al. Angiogenic properties of dehydrated human amnion/chorion allografts: therapeutic potential for soft tissue repair and regeneration. *Vascular Cell*. 2014. 6:10.

**Overall, it is vital to analyze the patient, the ability of the patient to abide by and follow the treatment plan, the wound, and the cause of the wound prior to utilizing conservative therapies.**

can prevent wounds and subsequent infections can make a significant difference in the success of amputation prevention programs.<sup>22</sup>

## Conclusions

Overall, it is vital to analyze the patient, the ability of the patient to abide by and follow the treatment plan, the wound, and the cause of the wound prior to utilizing conservative therapies. Though surgical intervention, such as amputations, in some cases may be the best course of treatment, various conservative therapies are available to the clinician to prevent this. Prevention of amputation by means of conservative therapy depends greatly on patient education and compliance, early diagnosis and treatment, and utilization of appropriate conservative therapies by the clinician. **PM**

## References

<sup>1</sup> Murdoch G, Wilson AB Jr, eds. Amputation: Surgical Practice and Patient Management. St Louis, Mo: Butterworth-Heinemann Medical; 1996.

<sup>2</sup> Tooms RE. Amputations. Crenshaw AH, ed. *Campbell's Operative Orthopedics*. 7th ed. St. Louis, Mo: Mosby-Year Book; 1987. Vol 1: 597-637.

<sup>3</sup> Lipsky BA, Berendt AR, Deery HG, et al. Diagnosis and treatment of diabetic foot infections. *Clin Infect Dis*. 2004 Oct 1. 39(7):885-910.

<sup>4</sup> Sheehan P, Edmonds M, Januzzi JL Jr, et al, for the Consensus Panel of the American Diabetes Association. Peripheral arterial disease in people with diabetes. *Diabetes Care*. 2003 Dec. 26(12):3333-41.

<sup>5</sup> National Diabetes Fact Sheet, ADA, 2004.

<sup>6</sup> Sender, Nicole L., and Manning, Linda,

31(8): 1679-1685.

<sup>9</sup> Lipsky, Benjamin A., et al. 2012 Infectious Diseases Society of America Clinical Practice Guideline for the Diagnosis and Treatment of Diabetic Foot Infections. *Clinical Infectious Disease*. 2015 June. 2012:54.

<sup>10</sup> Hingorani, Anil, et al. The management of the diabetic foot: A clinical practice guideline by the society of vascular surgery in collaboration with the American Podiatric medical association and the society for vascular medicine. *Journal of Vascular Surgery*. 2016. 63:3S-21S.

<sup>11</sup> E. K. Mooney, C. Lippitt, and J. Friedman, “Silver dressings (safety and efficacy reports),” *Plastic and Reconstructive Surgery*, vol. 117, no. 2, pp. 666-669, 2006.

<sup>12</sup> Wolcott Randy. Disrupting the bio-film matrix improves wound healing outcomes. *Journal of wound care*. 2015 Aug. 24(8):366-371.

<sup>13</sup> Heldin CH, Westermark B. Mechanism of action and in vivo role of platelet-derived growth factor. *Physiol Rev*. 1999;79:1283-1316.

<sup>14</sup> Mesut Mutluoglu, et al. Topical Oxygen for Chronic Wounds” A Pro/Con Debate. *J Am Coll Clin Wound Spec*. 2013 Dec; 5(3): 61-65.

<sup>15</sup> KHODR B., KHALIL Z. Modulation of inflammation by reactive oxygen species: implications for aging and tissue repair. *Free Radic Biol Med*. 2001;30:1-8.

<sup>16</sup> Gordillo G.M., Sen C.K. Revisiting the essential role of oxygen in wound healing. *Am J Surg*. 2003 Sep;186(3):259-263.

<sup>17</sup> Cronjé F.J. Oxygen therapy and wound healing—topical oxygen is not hyperbaric oxygen therapy. *S Afr Med J*. November 2005;95(11).

<sup>18</sup> Kubo M, Sonoda Y, Muramatsu R, Usui M: Immunogenicity of human amniotic membrane in experimental xenotransplantation. *Invest OphthalmolVis Sci* 2001, 42:1539-1546.

<sup>19</sup> Zelen CM, Serena TE, Denoziere G, Fetterolf DE: A prospective random-



**Dr. Priyanka Begur** received her Masters in Biomedical Sciences and a Doctor of Podiatric Medicine degree from Midwestern University in Glendale, Arizona in 2016. She then continued to complete a fellowship in Podiatric Research at the Carl T. Hayden Research Foundation. She is currently completing her residency at the Carl T. Hayden Veteran Affairs Medical Center in Phoenix, Arizona.



**Dr. Frykberg** received his Doctor of Podiatric Medicine degree from the California College of Podiatric Medicine in 1976. He completed his residency at the New England Deaconess Hospital/Harvard Medical School of Public Health with a concentration in quantitative methods. He is currently the Chief of Podiatric Medicine and Podiatry Residency Director at the Carl T. Hayden Veteran Affairs Medical Center in Phoenix, AZ. He is focused on treating high-risk foot problems and continues to do research and publish regarding diabetic foot ulcerations and the Charcot foot.